Effect of Tropolone, Azulene and Azulenequinone Derivatives on Prostaglandin E₂ Production by Activated Macrophage-like Cells

MASAYUKI NISHISHIRO¹, TERUO KURIHARA¹, HIDETSUGU WAKABAYASHI¹ and HIROSHI SAKAGAMI²

 ¹Department of Chemistry, Faculty of Science, Josai University, Sakado, Saitama;
²Division of Pharmacology, Department of Diagnostic and Therapeutic Sciences, Meikai University School of Dentistry, Sakado, Saitama, Japan

Abstract. We have previously reported that tropolone (T-3), 2,4-dibromo-7-methoxytropone (T-21), diethyl 2-chloroazulene-1,3-dicarboxylate (A-9),1,3-difluoroazulene (A-11),3-morpholino-1,5-azulenequinone (AQ-8) and 3,7-dibromo-1,5-azulenequinone (AQ-13) inhibited the nitric oxide (NO) production of lipopolysaccharide (LPS)-activated mouse macrophage-like RAW264.7 cells, with or without the inhibition of inducible NO synthase (iNOS) mRNA and protein expression. In order to confirm the anti-inflammatory potency, possible effects on prostaglandin (PG) E_2 production and the expression of enzymes involved in the arachidonic acid pathway were investigated. Among these six compounds, only A-9 effectively inhibited the PGE2 production of the LPS-stimulated RAW264.7 cells. Western blot analysis demonstrated that A-9 inhibited phospholipase A₂ (PLA₂), cyclooxygenase (COX)-2 and iNOS proteins only by 12, 45 and 42%, respectively. These data demonstrate the lack of correlation between the extent of inhibition of iNOS protein expression by tropolone or azulene derivatives and that of PGE₂, and suggest the possible antiinflammatory potency of A-9.

We have initiated the structure-activity study of tropolone and azulene derivatives. Among 107 compounds so far analyzed, tropolone (**T-3**), 2,4-dibromo-7-methoxytropone (**T-21**) (1), diethyl 2-chloroazulene-1,3-dicarboxylate (**A-9**), 1,3-difluoroazulene (**A-11**) (2), 3-morpholino-1,5-azulenequinone (**AQ-8**) and 3,7-dibromo-1,5-azulenequinone (**AQ-13**) (Figure 1) (3) inhibited the nitric oxide (NO) production of lipopolysaccharide (LPS)-activated mouse macrophage-like RAW264.7 cells (Table I), with (1, 3) or without (2) the

Correspondence to: Hidetsugu Wakabayashi, Faculty of Science, Josai University, Sakado, Saitama 350-0295, Japan. Tel: +81 049 2717959, Fax: +81 049 2717985, e-mail: hwaka@josai.ac.jp

Key Words: Diethyl 2-chloroazulene-1,3-dicarboxylate, RAW264.7 cells, macrophage, PGE₂, COX-2, PLA₂, anti-inflammatory effect.

inhibition of inducible NO synthase (iNOS) mRNA and protein expression. However, the extent of inhibition of iNOS expression differed considerably from compound to compound (1-3). Electron-spin resonance (ESR) spectroscopy showed that none of these compounds scavenged the NO released 1-hydroxy-2-oxo-3-(N-3-methyl-3-aminopropyl)-3from methyl-1-triazene (NO generator) in the presence of 2-(4carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxy1-3oxide (a spin trap agent) (1-3). In order to confirm the antiinflammatory potency of these six compounds, whether or not they inhibited the prostaglandin (PG) E2 production of LPSstimulated RAW264.7 cells was first investigated. In the arachidonic acid (AA) cascade, two enzymes are involved in the production of PGE₂. PG synthesis begins with the liberation of AA, the prime precursor, from membrane phospholipids by phospholipase A₂ (PLA₂). Subsequently, cyclooxygenase (COX) catalyzes the rate-limiting reactions for PG synthesis comprising the *bis*-cyclooxygenation of AA to form PGG₂ and the peroxidative reduction of this intermediate to PGH_2 (4). The cPLA₂ α is activated by proinflammatory cytokines or growth factors and catalyzes the AA from the cell surface membrane (5, 6). It is known that COX has two isoforms, COX-1 (constitutive isoform) and COX-2 (inducible isoform) (7). Therefore, the possible effects on the expression of these enzymes were also investigated.

Materials and Methods

Materials. The following chemicals and reagents were obtained from the indicated companies: Dulbecco's modified Eagle's medium (DMEM) (Invitrogen Corp, Carlsbad, CA, USA); fetal bovine serum (FBS) (Gemini Bio-Products, Woodland, CA, USA), LPS from *Escherichia coli* (Serotype 0111:B4) (Sigma Chemical Ind., St. Louis, MO, USA) and tropolone (**T-3**) (Tokyo Kasei Kogyo Co., Ltd., Tokyo, Japan).

Synthesis of tropolone, azulene and azulenequinone derivatives. **T-21** (8), **A-9** (9), **A-11** (9), **AQ-8** (10, 11) and **AQ-13** (10, 12, 13) were prepared, according to the reports in the references cited.





Tropolone (T-3)

Diethyl-2-chloroazulene-1,3 -dicarboxylate (A-9)



3-Morpholino-1,5azulenequinone (AQ-8)





2,4-Dibromo-7metoxytropone (T-21) 1,3-Difluoroazulene (A-11)



3,7-Dibromo-1,5azulenequinone (AQ-13)

Figure 1. Structure of compounds used in this study.

Cell culture. The RAW264.7 cells, established from the peritoneal fluid of BALB/c mice and showing the phenotypic characteristics of monocytes and macrophages (14), were subcultured in DMEM supplemented with 10% heat-inactivated FBS under a humidified 5% CO₂ atmosphere and incubated with various concentrations of test compounds in the presence or absence of LPS (100 ng/ml).

Assay for cytotoxic activity. The cytotoxic activity of the azulenes was determined by the MTT method, and expressed as absorbance at 540 nm of the MTT-stained cells. The 50% cytotoxic concentration (CC_{50}) was determined from the dose-response curve (1-3).

Assay for NO concentration. Near confluent RAW264.7 cells were incubated for 24 hours with each test sample in phenol red-free DMEM supplemented with 10% FBS, and the NO production was quantified by Greiss reagent, using the standard curve of NO_2^- . To eliminate the interaction between the sample and the Greiss reagent, the NO concentration in the culture medium without cells was also measured, and subtracted from the value obtained with the cells. The concentration that inhibited the LPS-stimulated NO production by 50% (50% inhibitory concentration: IC_{50}) was determined from the dose-response curve (1-3). The efficacy of inhibition of NO production was evaluated by the selectivity index (SI), which was calculated by the following equation: $SI=CC_{50} / IC_{50}$.

*Measurement of PGE*₂ *production*. RAW264.7 cells were subcultured in 24-well plates and incubated with various concentrations of test compounds in the presence of LPS (100 ng/ml). The culture supernatant was collected by centrifugation and the PGE₂ concentration was

determined by EIA kit (Cayman Chemical Co, Ann Arbor, MI, USA). Table I. Inhibition of NO production by LPS-stimulated RAW264.7 cells by tropolone, azulene and azulenequinone derivatives.

		Cytotoxic activity CC ₅₀ (µМ)		Inhibition of NO production IC ₅₀ (µM)		
Compound	MW	LPS(+)	LPS(-)	LPS(+)	SI	Ref
T-3	122	353	145	12	31	1
T-21	294	88	102	<1	>81	1
A-9	307	>261	>261	17	>15	2
A-11	164	368	353	12	30	2
AQ-8	243	381	315	12	26	3
AQ-13	316	179	165	8	22	3

Each value represents the mean from three independent experiments. CC_{50} : 50% cytotoxic concentration; IC_{50} : 50% inhibitory concentration; SI: selectivity index.

Western blotting. The cell pellets were lysed with 100 µl of lysis buffer (10 mM Tris-HCl [pH 7.6], 1% Triton[®] X-100, 150 mM NaCl, 5 mM EDTA-2Na and 2 mM phenylmethylsulfonyl fluoride (PMSF) for 10 minutes in ice water, and then incubated for 50 minutes at 4°C with RT-5 ROTATOR (Titec, Saitama, Japan). The cell lysates were centrifuged at 16,000 ×g for 20 minutes at 4°C to remove the insoluble materials and the supernatant was collected.



Figure 2. Effect of six selected compounds on PGE_2 production by LPS-activated RAW264.7 cells. RAW264.7 cells were incubated for 24 hours without (control), or with the indicated concentrations of tropolone (**T-3**), 2,4-dibromo-7-methoxytropone (**T-21**), diethyl 2-chloroazulene-1,3-dicarboxylate (**A-9**), 1,3-difluoroazulene (**A-11**), 3-morpholino-1,5-azulenequinone (**AQ-8**), or 3,7-dibromo-1,5-azulenequinone (**AQ-13**), and the extracellular concentration of PGE₂ was then measured.

The protein concentrations of the supernatant were measured by Protein Assay Kit (Bio Rad, Hercules, CA, USA). An equal amount of the protein from the cell lysates (10 µg) was mixed with 2x sodium dodecyl sulfate (SDS)-sample buffer (0.1 M Tris-HCl [pH 6.8], 20% glycerol, 4% SDS, 0.01% bromphenol blue, 1.2% 2mercaptoethanol), boiled for 10 minutes, applied to SDS-8% polyacrylamide gel electrophoresis and then transferred to polyvinylidene difluoride (PVDF) membrane. The membranes were blocked with 5% skim milk in phosphate-buffered saline (PBS(-)) plus 0.05% Tween 20 for 90 minutes and incubated for 90 minutes at room temperature with anti-PLA2, anti-COX-2 or anti-iNOS antibody (1:1,000; Santa Cruz Biotechnology, Delaware, CA, USA) or anti-actin antibody (1:2,000-4,000; Sigma) and then incubated with horseradish peroxidase-conjugated anti-mouse or goat IgG for 90 minutes at room temperature. The immunoblots were developed with a Western LightningTM Chemiluminescence Reagent Plus (Perkin Elmer Life Sciences, Boston, MA, USA) and analyzed on a Macintosh (Power Macintosh 7600/120) computer using the public domain NIH Image program (National Technical Information Service, Springfield, Virginia, USA, part number PB95-500195GEI).

Results

Among the six test compounds, only A-9 inhibited the PGE_2 production of the LPS-stimulated RAW264.7 cells (Figure 2). A-9 inhibited the PGE_2 production dose-dependently. A-9 at 2 or 20 μ M inhibited the PGE_2 production by 53.2 and 96.5%, respectively (Figure 3). Western blot analysis (Figure 4) demonstrated that A-9 inhibited PLA₂, COX-2 and iNOS proteins by only 12, 45 and 42%, respectively.

Discussion

The present study demonstrated that although tropolone (**T-3**), 2,4-dibromo-7-methoxytropone (**T-21**), 1,3-difluoroazulene (**A-11**), 3-morpholino-1,5-azulenequinone (**AQ-8**) and 3,7-



Figure 3. Dose response of A-9 in the inhibition of PGE_2 production by LPS-stimulated RAW264.7 cells. RAW264.7 cells were incubated for 24 hours without (control), or with the indicated concentrations of diethyl 2-chloroazulene-1,3-dicarboxylate (A-9), and the extracellular concentration of PGE_2 was then measured. Each value represents the mean and SD from three independent experiments.

dibromo-1,5-azulenequinone NO (AQ-13) inhibited production by LPS-stimulated mouse macrophage-like cells, these failed to inhibit the PGE₂ production of the same cells. This suggested that the inhibition of NO production by macrophages is not an appropriate parameter for evaluating anti-inflammatory activity. On the other hand, diethyl 2chloroazulene-1,3-dicarboxylate (A-9) inhibited PGE₂ production (IC₅₀<2 μ M) more efficiently than NO production $(IC_{50}=17 \ \mu M)$. These data suggested the importance of further investigating A-9 for its possible anti-inflammatory potential. It was unexpected that A-9 inhibited COX-2 protein expression only by 45%. It also inhibited the iNOS protein expression only partially and only marginally affected the PLA₂ protein expression. The binding assay with fluorescence-labeled LPS demonstrated that A-9 did not affect the binding of LPS to the cell surface receptor of the



Figure 4. Effect of diethyl 2-chloroazulene-1,3-dicarboxylate (A-9) on the intracellular concentration of PLA₂, iNOS and COX-2. RAW264.7 cells were incubated for 24 hours with the indicated concentrations of (A-9) in the absence (-) or presence (+) of 0.1 µg/ml LPS, and PLA₂ (A), COX-2 (B) or iNOS (C) protein levels were determined by western blot analysis. Each value represents mean and SD from three independent experiments.

RAW264.7 cells (Nishishiro, unpublished data), eliminating the possibility that the inhibitory effect of **A-9** was due to the inhibition of LPS binding to its receptor. It has been reported that PLA₂ α (15) and COX-2 (16) were activated by *S*-nitrosylation caused by iNOS. Therefore, it remains to be investigated whether **A-9** inhibits the nitrosylation of PLA₂ α and COX-2 proteins leading to the lower production of PGE₂.

We have previously reported that a lower concentration (16.3 μ M) of A-9 slightly reduced the viable cell number

of RAW264.7 cells with or without LPS stimulation (as shown by the dose-response curve in Figure 2) (2). This slight reduction of viability might overestimate the extent of the inhibition of NO production. However, such a minor growth inhibitory effect of A-9 at higher concentrations does not affect the inhibition of PGE₂ production observed at much lower concentration. Further study is required to elucidate the mechanism of the possible anti-inflammatory potency of A-9.

References

- 1 Yokoyama K, Hashiba K, Wakabayashi H, Hashimoto K, Satoh K, Kurihara T, Motohashi N and Sakagami H: Inhibition of LPSstimulated NO production in mouse macrophage-like cells by tropolones. Anticancer Res 24: 3917-3922, 2004.
- 2 Hashiba K, Yokoyama K, Wakabayashi H, Hashimoto K, Satoh K, Kurihara T, Motohashi N and Sakagami H: Inhibition of LPSstimulated NO production in mouse macrophage-like cells by azulenes. Anticancer Res 24: 3939-3944, 2004.
- 3 Nishishiro M, Arikawa S, Wakabayashi H, Hashimoto K, Satoh K, Yokoyama K, Unten S, Kakuta H, Kurihara T, Motohashi N and Sakagami H: Inhibition of LPS-stimulated NO production in mouse macrophage-like cells by azulenequinones. Anticancer Res 25: 4157-4164, 2005.
- 4 Odau S, Gabler C, Holder C and Einspanier R: Differential expression of cyclooxygenase 1 and cyclooxygenase 2 in the bovine oviduct. J Endocrinol *191*: 263-274, 2006.
- 5 Kudo I and Murakami M: Phospholipase A₂ enzymes. Prostaglandins Other Lipid Mediat *68-69*: 3-58, 2002.
- 6 Ghosh M, Tucker DE, Burchett SA and Leslie CC: Properties of the Group IV phospholipase A₂ family. Prog Lipid Res 45: 487-510, 2006.
- 7 O'Banion MK, Winn VD and Young DA: cDNA cloning and functional activity of a glucocorticoid-regulated inflammatory cyclooxygenase. Proc Natl Acad Sci *89*: 4888-4892, 1992.
- 8 Wakabayashi H, Yokoyama K, Hashiba K, Hashimoto K, Kikuchi H, Nishikawa H, Kurihara T, Satoh K, Shioda S, Muto S, Terakubo S, Nakashima H, Motohashi N and Sakagami H: Cytotoxic activity of tropolones against human oral tumor cell lines. Anticancer Res 23: 4757-4764, 2003.
- 9 Wakabayashi H, Hashiba K, Yokoyama K, Hashimoto K, Kikuchi H, Nishikawa H, Kurihara T, Satoh K, Shioda S, Saito S, Kusano S, Nakashima H, Motohashi N and Sakagami H: Cytotoxic activity of azulenes against human oral tumor cell lines. Anticancer Res 23: 4747-4756, 2003.

- 10 Nozoe T and Takeshita H: Chemistsry of azulenequinones and their analogues. Bull Chem Soc Jpn 69: 1149-1178, 1996.
- 11 Nozoe T, Wakabayashi H, Shindo K and Ishikawa S: Preparation of various azulenequinone derivatives by a nucleophilic substitution of 3-bromoazulenequinone synthones. Chem Lett, pp. 27-28, 1995.
- 12 Wakabayashi H, Kurihara T, Shindo K, Tsukada M, Yang P-W, Yasunami M and Nozoe T: Synthesis and reactivities of 1,5- and 1,7-azulenequinone derivatives. A highly convenient, one-pot synthesis of 3-bromo-1,5- and -1,7-azulenequinones by bromination of azulene. J Chin Chem Soc 45: 391-400, 1998.
- 13 Nozoe T, Wakabayashi H, Shindo K, Kurihara T, Ishikawa S and Kageyama M: A highly convenient, one-pot synthesis of 3-bromo-1,5- and -1,7-azulenequinones by polybromination of azulene. Chem Lett, pp. 25-26, 1995.
- 14 Ralph R and Nakoinz I: Antibody-dependent killing of erythrocyte and tumor targets by macrophage-related cell lines: enhancement by PPD and LPS. J Immunol *119*: 950-954, 1977.
- 15 Xu L, Han C, Lim K and Wu T: Activation of cytosolic phospholipase A2α through nitric oxide-induced *S*-nitrosylation: involvement of inducible nitric oxide synthase and cyclooxygenase-2. J Biol Chem 283: 3077-3087, 2007.
- 16 Kim SF, Huri DA and Snyder SH: Inducible nitric oxide synthase binds, *S*-nitrosylates, and activates cyclooxygenase-2. Science *310*: 1966-1970, 2005.

Received May 19, 2008 Revised November 20, 2008 Accepted December 1, 2008